

Supplementary Appendix

This appendix has been provided by the authors to give readers additional information about their work.

Supplement to: Monti MM, Vanhaudenhuyse A, Coleman MR, et al. Willful modulation of brain activity in disorders of consciousness. *N Engl J Med* 2010. DOI: [10.1056/NEJMoa0905370](https://doi.org/10.1056/NEJMoa0905370).

Online Appendix for:

Willful modulation of brain activity in disorders of consciousness

Martin M. Monti, Ph.D., Audrey Vanhaudenhuyse, M.Sc., Martin R. Coleman, Ph.D.,
Melanie Boly, M.D., John D. Pickard, F.R.S.C., F.Med.Sci., Luaba Tshibanda, M.D.,
Adrian M. Owen, Ph.D., and Steven Laureys, M.D., Ph.D.

This document includes:

1. Inclusion Criteria
2. fMRI Data Acquisition Parameters and Pre-Processing
3. Relative Similarity Metric
4. Healthy Volunteer Results
5. Similarity Analysis Result for the Patient
6. Voluntary vs. Automatic Brain Processing

References

1. Inclusion Criteria

Both the Cambridge and the Liège sites routinely admit brain injury patients (VS and MCS) for evaluation with fMRI, from a network of referring centers. In both locations, patients are neither pre-selected nor pre-screened on the basis of bedside examinations. The main constraint to admitting a patient is where they require paramagnetic medical apparatus that may not be suitable for entering the fMRI environment. In addition, in Cambridge, patients that appear clearly hyperkinetic and unlikely to remain sufficiently still throughout the imaging session are, when such a situation is evident, not admitted. In Liege, patients undergo structural scanning under sedation. Many of these patients, however, during the functional scans (when they are not sedated) exhibit excessive movements (several centimeter), rendering the data not analyzable.

2. fMRI Data Acquisition Parameters and Pre-Processing

fMRI Data Acquisition. Volunteer data was collected at the MRC Cognition and Brain Sciences Unit, Cambridge (UK) on a 3T Tim Trio Siemens system. Patient data was collected at the Wolfson Brain Imaging Centre, Cambridge, on a 3T Siemens Tim Trio and a 3T Brucker system, and at the Liège University Hospital (Belgium) on a 3T Siemens Allegra. T1-weighted images were acquired with a 3D MP-RAGE sequence (TR 2300 ms, TE 2.47 ms, TI 900 ms, 150 slices, 1x1x1.2 mm resolution). T2* sensitive images were acquired using an echo planar sequence (TR 2000 ms, TE 30 ms, 32 descending axial slices, 3x3x3.75 mm resolution on the Siemens machines, and TR 1100 ms, TE 27.5 ms, 21 interleaved transverse slices, 4 mm thickness on the Bruker system).

fMRI Data pre-processing. Analysis methods were performed using FSL 4.1 (FMRIB Software Library, Oxford University).¹ Prior to functional analyses, a series of pre-processing steps were performed. First, signal from extraneous non-brain tissue was removed using BET (Brain Extraction Tool).² Each individual echo planar imaging (EPI) time-series was motion corrected to the middle time point using a 6 parameter, rigid-body method (as implemented in MCFLIRT).³ Data were then band-pass filtered (2.8 – 60 s) and smoothed using a Gaussian kernel of 5 mm FWHM. Autocorrelation was corrected with a pre-whitening technique (as implemented in FEAT; fMRI Expert Analysis Tool).⁴

3. Relative Similarity Metric

Similarity of brain activation between question and localizer scans was assessed according to the Euclidean distance. Specifically, activity in each scan was re-described as a point within a two dimensional plane with axes corresponding to the activation seen in each ROI (SMA, PPA). If one defines *total distance* as the sum of the distances separating a given question scan from the two localizers, the *relative similarity* (rs) between a given question and each localizer is equal to one minus the ratio of the distance between the question and each localizer, and the total distance. For example, the relative similarity of a given question Q_i to each localizer (tennis localizer, TL; and navigation localizer, NL) can be obtained as follows (with $d(x,y)$ representing the Euclidean distance separating point x from point y):

$$rs(Q_i, TL) = 1 - \left(\frac{d(Q_i, TL)}{d(Q_i, TL) + d(Q_i, NL)} \right)$$

$$rs(Q_i, NL) = 1 - \left(\frac{d(Q_i, NL)}{d(Q_i, TL) + d(Q_i, NL)} \right)$$

In two-dimensional space, the smaller the distance separating a question and a localizer scan, the greater the relative similarity.

4. Healthy Volunteer Results

Analysis of the localizer data for each healthy volunteer revealed two consistent patterns of activity in response to motor and spatial imagery. For 7 out of 16 volunteers, each ROI selectively responded to just one type of imagery, with the SMA responding to motor imagery only and PPA responding to spatial imagery only. For the remaining 9 volunteers, motor imagery activated the SMA alone, but spatial imagery activated both the PPA and, to a lesser extent, the SMA. Noticeably, whichever pattern was detected in the localizers, that same pattern was observed during the question scans. The similarity analysis successfully 'decoded', with 100% accuracy, the answer provided by modulation of brain activity alone to each of the 48 questions (3 questions per subject). Indeed, the pattern of ROI activation in each question scan was always more similar to the imagery task associated with the factually correct answer (see Table A1).

Table A1. Relative similarity data for 16 healthy volunteers. (In bold the imagery task that corresponded, for each question, to the correct answer.)

		% Similarity to Localizers		
		Question 1	Question 2	Question 3
sub 1	Motor Imagery Localizer	84.65	14.94	14.96
	Spatial Imagery Localizer	15.35	85.06	85.04
sub 2	Motor Imagery Localizer	94.23	25.89	81.03
	Spatial Imagery Localizer	5.77	74.11	18.97
sub 3	Motor Imagery Localizer	80.01	29.92	82.30
	Spatial Imagery Localizer	19.99	70.08	17.70
sub 4	Motor Imagery Localizer	22.32	88.83	81.76
	Spatial Imagery Localizer	77.68	11.17	18.24
sub 5	Motor Imagery Localizer	27.26	84.73	64.93
	Spatial Imagery Localizer	72.74	15.27	35.07
sub 6	Motor Imagery Localizer	12.83	81.66	23.42
	Spatial Imagery Localizer	87.17	18.34	76.58
sub 7	Motor Imagery Localizer	82.84	38.66	30.56
	Spatial Imagery Localizer	17.16	61.34	69.44
sub 8	Motor Imagery Localizer	25.92	13.18	86.72
	Spatial Imagery Localizer	74.08	86.82	13.28
sub 9	Motor Imagery Localizer	14.94	65.39	11.83
	Spatial Imagery Localizer	85.06	34.61	88.17
sub 10	Motor Imagery Localizer	87.17	77.99	13.02
	Spatial Imagery Localizer	12.83	22.01	86.98
sub 11	Motor Imagery Localizer	85.50	88.33	5.68
	Spatial Imagery Localizer	14.50	11.67	94.32
sub 12	Motor Imagery Localizer	84.26	17.92	87.28
	Spatial Imagery Localizer	15.74	82.08	12.72
sub 13	Motor Imagery Localizer	83.76	17.87	17.27
	Spatial Imagery Localizer	16.24	82.13	82.73
sub 14	Motor Imagery Localizer	92.84	12.86	8.61
	Spatial Imagery Localizer	7.16	87.14	91.39
sub 15	Motor Imagery Localizer	96.92	11.11	38.42
	Spatial Imagery Localizer	3.08	88.89	61.58
sub 16	Motor Imagery Localizer	78.64	12.12	67.44
	Spatial Imagery Localizer	21.36	87.88	32.56

5. Similarity Analysis Results for the Patient.**Table A2.** Relative similarity data for the patient. (In bold the imagery task that corresponded, for each question, to the correct answer.)

		% Similarity to Localizers					
		Question 1	Question 2	Question 3	Question 4	Question 5	Question 6
Patient	Motor Imagery Localizer	33.93	24.01	82.31	66.89	24.88	51.93
	Spatial Imagery Localizer	66.07	75.98	17.69	33.11	75.12	48.07

6. Voluntary vs. Automatic Brain Responses

Is there any possibility that this patient was not conscious, yet able to generate appropriate answers to autobiographical questions 'automatically' in response to the questions? Recent evidence suggests that single words can, under certain circumstances, elicit wholly automatic neural responses in the absence of conscious awareness.

However, such responses last for a few seconds at most and, unsurprisingly, occur in regions of the brain that are associated with word processing.⁵ In contrast, the responses in the patient presented here were sustained across the 30 sec epochs in the absence of any further stimulation and were observed in regions that are known to be involved in the two imagery tasks.^{6,7} More importantly, in the current study, the same neutral word ('*answer*') was used to cue a response, irrespective of which imagery task was to be performed. This precludes any possibility that the observed activity occurred automatically (i.e. in the absence of awareness) since in different questions an identical cue yielded different, yet predicted, BOLD responses. These responses could, therefore, only depend on the patient's conscious decision (or 'mindset') about which answer to give (see also ref. 8 for discussion).

With respect to the novel communication method presented in the main text, in order to 'answer' a question, the patient was first required to select which of the two imagery tasks was appropriate for the answer that he intended to give ('yes' or 'no') and to engage in that type of imagery when cued with the word '*answer*' and disengage (or relax) when cued by the word '*relax*.' Each period of imagery required his sustained involvement in the task in order to generate continuous activity in the target ROI across each 30 second epoch. Moreover, in order for a statistically reliable '*answer*' to be detectable he was required to repeat each imagery task 5 times (per question).

Sustained, time-locked and repeated activity within well characterized neuroanatomical regions requires a level of cognitive processing that includes language comprehension, memory, attention and voluntary or 'willful' behavior.

References

1. Smith SM, Jenkinson M, Woolrich MW, et al. Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage* 2004;23:208-219.
2. Smith SM. Fast robust automated brain extraction. *Hum. Brain Mapp.* 2002;17:143-155.
3. Jenkinson M, Bannister PR, Brady JM, Smith SM. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *NeuroImage* 2002;17:825-841.
4. Woolrich MW, Ripley BD, Brady JM, Smith SM. Temporal autocorrelation in univariate linear modelling of fMRI data. *NeuroImage* 2001;14:1370-1386.
5. Hauk O, Johnsrude I, Pulvermüller F. Somatotopic representation of action words in human motor and premotor cortex. *Neuron* 2004;41:301-7.
6. Boly M, Coleman MR, Hampshire A *et al.* When thoughts become action: an fMRI paradigm to study volatile brain activity in non-communicative brain injured patients. *Neuroimage* 2007;36:979-92.
7. Weiskopf N, Mathiak K, Bock SW *et al.* Principles of a brain-computer interface (BCI) based on real-time functional magnetic resonance imaging (fMRI). *IEEE Trans Biomed Eng* 2004;51:966-70.
8. Owen AM, Coleman MR, Boly M *et al.* Response to Comments on "Detecting Awareness in the Vegetative State". *Science* 2007; 315: 1221.